

LIVER DISEASE BURDEN AND CLINICAL FOLLOW-UP DURING A LIVER HEALTH PROMOTION INTERVENTION INTEGRATING NON-INVASIVE LIVER DISEASE SCREENING IN DRUG AND ALCOHOL SETTINGS: THE LIVERLIFE STUDY

Grebely J¹, Marshall AD¹, Krahe M², Erratt A¹, Telenta J³, Treloar C⁴, Jones SC³, Bath N⁶, How-Chow D⁷, Byrne J⁸, Harvey P⁹, Dunlop A^{10,11}, Applegate TL¹, Lamoury F¹, Mowat Y¹, Jauncey M¹², Read P^{1,13}, Gilliver R¹³, Smith J¹⁴, Collie T¹⁵, and Dore GJ¹

¹The Kirby Institute, UNSW Australia, NSW Australia; ²Menzies Health Institute Queensland, Griffith University, QLD Australia; ³Centre for Health and Social Research, Australian Catholic University, VIC Australia; ⁴Centre for Social Research in Health, UNSW Australia, NSW Australia; ⁵NSW Users and AIDS Association, Inc., NSW Australia; ⁶NSW Health, ⁷St Vincent's Hospital Sydney, NSW Australia; ⁸Australian Injecting and Illicit Drug Users League, ACT Australia; ⁹Hepatitis NSW, Australia; ¹⁰University of Newcastle, Newcastle, NSW, Australia; ¹¹Drug and Alcohol Clinical Services, Hunter New England Local Health District, Newcastle, NSW, Australia; ¹²Sydney Medically Supervised Injecting Centre, NSW, Australia; ¹³Kirketon Road Centre, NSW Australia; ¹⁴Matthew Talbot Hostel, St Vincent de Paul Society NSW Support Services, NSW Australia; ¹⁵Coffs Harbour Drug and Alcohol Service, NSW, Australia

Background: Liver disease burden among people who inject drugs (PWID) is high, yet few strategies to enhance liver disease screening have been evaluated. The aim of this study was to assess factors associated with severe fibrosis/cirrhosis and follow-up among PWID participating in a liver health promotion campaign.

Methods: The LiveRLife campaign involved: 1) educational resource development; 2) resource testing; and 3) implementation. Between May-October 2014, participants were enrolled in an observational cohort study with recruitment from four clinics in Australia (one community-based primary health care clinic, two opioid substitution treatment clinics, and one medically supervised injecting centre). Participants received educational material, clinical assessment, transient elastography (TE) assessment, dried blood spot testing, and completed a knowledge survey. Qualitative HCV RNA testing was performed on dried blood spots collected from the study.

Results: Of 253 participants (mean age=43), 68% were male, 71% had injected in the past month, and 68% were HCV RNA+. Overall, 68% had no/mild fibrosis (F0/F1, ≥ 2.5 - ≤ 7.4 kPa), 13% moderate fibrosis (F2, ≥ 7.5 - ≤ 9.4 kPa), 10% severe fibrosis (F3, ≥ 9.5 - ≤ 12.4 kPa), and 9% had cirrhosis (F4, ≥ 12.5 kPa). The proportion of people with severe fibrosis/cirrhosis (F3/F4, 19%) was higher in those who were >50 years (33% vs. 15%, $P=0.003$), male (23% vs. 11%, $P=0.069$), and those HCV RNA+ (24% vs. 10%, $P=0.011$). In adjusted analysis, age >50 years (OR 2.91, 95%CI, 1.42, 5.95) and being HCV RNA+ (OR 2.61, 95%CI, 1.08, 6.28) were associated with severe fibrosis/cirrhosis (F3/F4). Sixty percent (n=152) returned for a follow-up nurse/specialist assessment.

Conclusion: Liver disease burden in this population was high and was associated with age >50 years and HCV RNA positive status. The majority of people assessed for liver disease returned for a follow-up assessment by a nurse/specialist, supporting the inclusion of TE in HCV-related care.

Disclosure of Interest Statement: The study was supported in part by a research grant from Investigator-Initiated Studies Program of Merck Sharp & Dohme Corp. The opinions expressed in this paper are those of the authors and do not necessarily represent those of Merck Sharp & Dohme Corp. The Kirby Institute is funded by the Australian Government Department of Health and Ageing. The views expressed in this publication do not necessarily represent the position of the Australian Government. GD is supported by a National Health and Medical Research Council Practitioner Research Fellowships. JG is supported by a National Health and Medical Research Council Career Development Fellowship. SJ is supported by an Australian Research Council Future Fellowship.